

Iris-Claw Intraocular Lens (Artisan IOL) Implantation: Indications

BY SARA AHMED AND EAMONN O'CONNELL

CPD record / Reader knowledge check

Please print out our form, complete the questions below and sign.
Keep for your records.



1. Which of the following was the most common indication for Artisan IOL implantation in this study?
 - ☐ Subluxated intraocular lens
 - ☐ Dislocated intraocular lens
 - ☐ Aphakia
 - ☐ Opacified intraocular lens
2. Which complication showed a higher incidence specifically in retro-pupillary implantation?
 - ☐ Corneal decompensation
 - ☐ Cystoid macular oedema (CMO)
 - ☐ Raised intraocular pressure (IOP)
 - ☐ Iris transillumination defects
3. What was the most common postoperative complication overall?
 - ☐ Corneal decompensation
 - ☐ Iris transillumination defects
 - ☐ Cystoid macular oedema (CMO)
 - ☐ Lens dislocation
4. Which statement best describes the visual outcomes reported in this study?
 - ☐ Visual outcomes were significantly worse in patients with comorbidities
 - ☐ BCVA was significantly better in anterior chamber implantation
 - ☐ No association was found between visual outcomes and comorbidities or lens position
 - ☐ Most patients had BCVA worse than 6/60
5. What was the primary purpose of performing surgical peripheral iridotomy (PI) in these patients?
 - ☐ Improve visual acuity outcomes
 - ☐ Reduce risk of retinal detachment
 - ☐ Reduce risk of postoperative high IOP/pupillary block
 - ☐ Prevent cystoid macular oedema
6. Which of the following was the most common pre-existing ocular comorbidity?
 - ☐ Pseudo exfoliation syndrome
 - ☐ Marfan syndrome
 - ☐ Previous retinal detachment repair
 - ☐ Glaucoma



eyenews®

